

## OSMOREGULATION IN THE BROOK TROUT, *SALVELINUS FONTINALIS*—I. DIEL, PHOTOPERIOD AND GROWTH RELATED PHYSIOLOGICAL CHANGES IN FRESHWATER

STEPHEN D. MCCORMICK\* and ROBERT J. NAIMAN

Woods Hole Oceanographic Institution, Woods Hole, Massachusetts 02543, USA

(Received 23 January 1984)

**Abstract**—1. Brook trout (*Salvelinus fontinalis*) raised from eggs under two photoperiod and two feeding regimes were tested for physiological changes preparatory for transition from freshwater to seawater. Size, age, growth rate, photoperiod, and diel rhythms were examined for possible influences on plasma osmolarity,  $[Na^+]$ ,  $[Cl^-]$ ,  $[K^+]$ ,  $[Mg^{2+}]$ , thyroxine concentration, hematocrit, and gill  $Na^+$ ,  $K^+$ -ATPase activity of brook trout in freshwater.

2. Significant diel cycles were found in plasma osmolarity,  $[Na^+]$  and thyroxine concentration.

3. Significant size and/or age related changes occurred for plasma osmolarity,  $[Na^+]$ ,  $[K^+]$  and hematocrit, but could explain little of their total variation ( $0.02 < r^2 < 0.18$ ).

4. A sexually dimorphic response to photoperiod was observed in hematocrit for both mature and immature fish, with hematocrit of mature females declining in autumn and hematocrit of immature males increasing in autumn.

5. Gill  $Na^+$ ,  $K^+$ -ATPase activity did not respond to photoperiod or feeding treatment and showed no change with size or age.

6. Plasma thyroxine levels responded to feeding and photoperiod treatment. There was a significant correlation between the percent mean difference in plasma thyroxine and the mean difference in growth rate between high and low feed fish ( $r^2 = 0.51$ ), suggesting a relationship between thyroxine and growth.

### INTRODUCTION

Anadromous salmonids are truly euryhaline over only a limited portion of their life cycle. A variety of physiological and morphological changes associated with transformation from freshwater parr to the migratory smolt occur wholly in freshwater (Gorbman *et al.*, 1982) and are presumably adaptive to the fishes existence in seawater. These changes, which include guanine deposition in skin and scale, increased hypoosmoregulatory ability, increased activity of gill  $Na^+$ ,  $K^+$ -ATPase and a surge of plasma thyroxine, are responsive to internal changes such as size and growth (Wedemeyer *et al.*, 1980) and environmental cues such as lunar and seasonal rhythms (Grau *et al.*, 1982). While smoltification occurs in migratory species of *Oncorhynchus* and *Salmo*, little is known regarding adaptations to euryhalinity in the charrs, which comprise the salmonid genus *Salvelinus*. This genus is regarded as a primitive group relative to *Salmo* and *Oncorhynchus* and displays a more generalized and restricted pattern of seaward migration (Hoar, 1976).

Although most populations of the brook trout (*Salvelinus fontinalis*) are restricted to freshwater, many coastal rivers in northeastern North America contain anadromous brook trout. In northern latitudes brook trout migrations are characterized by spring emigrations and coastal sea residence which

lasts for 2-4 months (White, 1940; Wilder, 1952; Dutil and Power, 1980; Castonguay *et al.*, 1982). In the southern portion of its range migrations are more variable, often occurring in the fall (Mullan, 1958). Sea ranching experiments resulted in spring emigrations and fall returns in which migrating fish obtained growth rates 4 to 5 greater than those of fish that remained in fresh water, and returned at rates between 30% and 60% (Whoriskey *et al.*, 1981).

The studies reported here were designed to reveal the factors which limit anadromy in natural populations of brook trout, and which would affect the culture and stocking of brook trout for natural enhancement, sea ranching and farming. In addition, the physiological adaptations of a less specialized anadromous species such as brook trout represent a "primitive archetype" from which we can interpret specialization made by more advanced salmonids. In conjunction with a study of factors which limit salinity tolerance and hypoosmoregulatory ability in brook trout (McCormick and Naiman, 1984a), we have investigated the preparatory physiological adaptations for euryhalinity that are characteristic of smolting salmonids which may occur in the facultatively anadromous brook trout. Specifically, we report here the effect of size, age, growth and photoperiod on gill  $Na^+$ ,  $K^+$ -ATPase and plasma thyroxine levels of brook trout in freshwater. Plasma osmolarity,  $[Na^+]$ ,  $[Cl^-]$ ,  $[K^+]$  and  $[Mg^{2+}]$ , and hematocrit were also measured to determine possible ontogenetic changes in hyperosmoregulation which may signal a physiological change relating to preparatory adaptations for hypoosmoregulation. Diel cycles of blood parameters were examined to verify

\*Present Address: Fisheries and Environmental Sciences, Department of Fisheries and Oceans, Biological Station, St. Andrews, N.B. E0G 2X0 Canada.




Fig. 1. Experimental design and fish culture conditions consisting of two photoperiod regimes (one normal, one 3-month delayed) and two feeding regimens (high feed and low feed).

their existence as reported for other teleosts (Hannah and Pickford, 1981; Eales *et al.*, 1981; Spieler and Noeske, 1979; Osborn *et al.*, 1978), and to analyze their possible effects in determining seasonal cycles.

#### MATERIALS AND METHODS

##### Experimental animals

20,000 fertilized brook trout eggs were obtained from the Massachusetts State Fish Hatchery at Sandwich. Although the breeding stock has been exclusively freshwater for the last 30 years (Lloyd Raymond, Hatchery Manager, personal communication), studies indicate the strain displays anadromy similar to wild stocks when released into coastal rivers (Mullan, 1958). Fertilized eggs were transported to the Woods Hole Oceanographic Institution's Shore Lab facility and supplied with 10–11°C well water. Eggs were randomly divided into two annually cycling photoperiods corresponding to a latitude of 42°N; one photoperiod cycle corresponded to the normal calendar date (longest day June 21, shortest day December 21), while the other was 3 month delayed from the norm (longest day September 21, shortest day March 21, Fig. 1). Daylength was changed every 5 days. Sunrise and sunset were simulated each day by a 15 min period of gradual illumination or dimming of incandescent bulbs. Beginning and end of daylight period were delayed 2 hr from Eastern Standard Time. Vita-Lite spiralux fluorescent bulbs and incandescent bulbs were used to simulate daylight.

Eggs were hatched in 250 l flow-through hatchery troughs. After first feeding fish were transferred to 1000 l flow-through tanks which received supplemental aeration. Within one week after feeding fish were divided randomly, within each photoperiod treatment, into two feeding groups. For 4 weeks, after first feeding, fish in each group were fed equal amounts. The high feed group was fed "maximum" rations (percent body weight of feed per day decreased with increasing body size; Leitritz and Lewis, 1976). The low feed group was fed approximately half the amount, per unit body weight, fed the high feed group. Each group was fed 4 to 5 times daily during daylight hours for the first several months after hatching, 1 to 3 times daily thereafter. Some fish, which were used as control fish in a related study of salinity tolerance, were maintained for 3–6 weeks in 100 l flow-through tanks prior to sampling. Feeding behavior of fish in these tanks was attenuated for a period of 4–5 days after transfer and then returned to normal.

Every 6–8 weeks, at least 25% of the fish from a 1000 l tank in each feeding group in the normal photoperiod were weighed. Fish were dip-netted, anesthetized, blot-dried on a moist chamois cloth, fork length was measured to the nearest mm and fish were weighed to the nearest 0.01 g.

Specific growth rates ( $G_w$ ) were calculated using the following formula:

$$G_w = \left( \frac{\log_e W_t - \log_e W_0}{t} \right) \cdot 100$$

where  $W_t$  is the weight at time  $t$ ,  $W_0$  is the weight at time 0, and  $t$  is time in days. In order to compare growth rates of animals of different sizes, the  $\log_e G_w$  of a fish of unit size was calculated (Jobling, 1983) using the following equation:

$$a = \log_e G_w + b \log_e W_t$$

where  $a$  is the  $\log_e G_w$  of a fish of unit size and  $b$  is the slope of the linear relationship between  $\log_e G_w$  and  $\log_e W_t$ . An experimentally derived value of  $b = -0.47$  was used in all calculations of  $a$ . A generalized value of  $b = -0.41$  for salmonids was reported by Brett (1979).

##### Blood and gill tissue sampling

Fish were starved overnight prior to blood collection which occurred between one hour after dawn and one hour prior to sunset. Brook trout were removed from tanks, placed in aerated transfer buckets for <2 min, and transferred immediately to a 0.4 ml/l phenoxyethanol-water solution for 30–60 s. Anesthetized animals were blotted with a damp chamois cloth, fork length was measured and fish were weighed. After severing of the caudal peduncle, blood from each fish was collected from the dorsal aorta into two heparinized micro-hematocrit tubes, which were then sealed at one end with vinyl plastic putty. Hematocrit tubes were centrifuged for 5 min at 5500 rpm, hematocrits (% red blood cells) read, and plasma removed for later analysis (see below). Gill arches were removed immediately after blood collection for gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity, and gonads were removed and weighed.

##### Analytical techniques

Plasma was withdrawn from hematocrit tubes with a positive displacement Hamilton syringe. Osmolarity and  $[\text{Cl}^-]$  were measured immediately with a Wescor Vapor Pressure Osmometer and Buchler-Cotlove Chloridometer, respectively. A 5  $\mu\text{l}$  plasma sample was diluted in 495  $\mu\text{l}$  deionized water in a 2.5 ml acid-washed, polyethylene vial and stored in the dark for a maximum of 48 hr. Reference standards were diluted and stored in the same manner.  $[\text{Na}^+]$  and  $[\text{K}^+]$  were measured using flame emission spectrophotometry;  $[\text{Mg}^{2+}]$  was measured using atomic absorption spectrophotometry. An additional dilution of 1:250 was made in duplicate vials for measuring  $[\text{Na}^+]$ . It was found that  $\text{Na}^+$  had a small but detectable interference with  $\text{K}^+$  that was constant over the physiological range of plasma  $[\text{Na}^+]$ . To correct for this error, 150 mmole  $[\text{Na}^+]/\text{l}$

was added to all  $[K^+]$  standards. Intraassay coefficients of variation, including dilutions, were 0.8%, 0.6%, 2.0%, 1.5% and 1.0% ( $n = 5$ ) for osmolarity,  $[Cl^-]$ ,  $[Na^+]$ ,  $[K^+]$  and  $[Mg^{2+}]$ , respectively. Interassay coefficients of variation for  $[Na^+]$ ,  $[K^+]$  and  $[Mg^{2+}]$  were 2.0% ( $n = 8$ ), 3.8% ( $n = 7$ ) and 2.8% ( $n = 9$ ), respectively.

For thyroxine determination 25 to 40  $\mu$ l of plasma was removed from hematocrit tubes, placed in a 250  $\mu$ l polyethylene microcentrifuge vial and stored at  $-17^\circ C$  for up to 6 months. For fish less than 8.5 cm fork length (high feed group in June 1981 and low feed group from June 1 to August 11, 1981), plasma from two to four fish was pooled. For statistical analysis the value of a pooled sample was treated as that of a single fish (e.g. for  $n = 5$ , up to 20 fish were actually used). Samples were thawed and duplicate 10  $\mu$ l samples withdrawn and analyzed using a competitive binding radioimmunoassay (Dickhoff *et al.*, 1978). Charcoal stripped brook trout plasma was used to make all standards. Sensitivity of the thyroxine radioimmunoassay was approximately 0.25 ng/ml. Intraassay variation was  $\pm 10\%$  ( $n = 5$ ), interassay variation was  $\pm 13\%$  ( $n = 3$ , with four replicates each).

Primary gill filaments (0.05–0.2 g wet weight) were trimmed from ceratobranchials and stored in 1 ml Sucrose-EDTA-imidazole (SEI) solution (0.3 mmole/l sucrose, 0.02 mmole/l disodium ethylenediamine tetraacetate and 0.1 mmole/l imidazole adjusted to a final pH of 7.1 with HCl), at  $-17^\circ C$ . Gill  $Na^+$ ,  $K^+$ -ATPase activity was determined by the method of Zaugg (1982). Protein determinations were done according to Lowry *et al.* (1951) as modified by Miller (1959) using bovine serum albumin as standard. Gill  $Na^+$ ,  $K^+$ -ATPase activity is expressed as  $\mu$ M inorganic phosphate per mg protein per hr ( $\mu$ MP<sub>i</sub>·mg prot.<sup>-1</sup>·hr<sup>-1</sup>). To determine the reproducibility of the assay and assess the effect of storage, primary filaments from several fish were pooled and then separated into vials containing 1 ml SEI solution. Intra- and interassay coefficients of variation were 7% ( $n = 6$ ) and 21% ( $n = 4$ , with five replicates each), respectively. The mean activities of five samples decreased 9.6% after 40 days and an additional 13.8% after 80 days. Maximum storage time of gill tissue was 50 days.

#### Statistical analysis

The experimental design (Fig. 1) is a  $2 \times 2$  matrix, such that feeding treatment, which affects growth and size simultaneously, is tested twice (once in normal photoperiod, once in 3-month delayed photoperiod), and photoperiod treatment is tested twice (once on high feed fish, once on low feed fish). To analyze our results, four separate two-way analyses of variance for each physiological variable were used to assess the effect of feeding and photoperiod treatment. Data from each day were entered in columns and two feeding or photoperiod treatments entered as rows. A photoperiod or feeding effect was deemed significant when the group (row) effect was  $P < 0.05$ . This method avoided the false influence of day-to-day variations in fish response to uncontrolled stimuli and day-to-day variations in analytical techniques.

## RESULTS AND DISCUSSION

### Size and growth rate

Length and weight of fish in high and low feed groups under normal photoperiod conditions are shown in Fig. 2(A,B) and were similar for high and low feed groups, respectively, in 3-month delayed photoperiod. Growth rates in each feeding treatment (Fig. 2(C)) were similar at low body weight (shortly after first feeding), were lower in the low feeding group at intermediate body weight, then became similar at higher body weights. Growth rate per unit




Fig. 2. Effect of feeding treatment on size and growth rate. (A) Length and (B) weight of normal photoperiod fish in high feed (squares) and low feed (circles) groups as a function of time. Mean value of 40–75 fish per sampling date. (C) Log<sub>e</sub> specific growth rate ( $G_w$ ) as a function of the natural logarithm of fish weight ( $g$ ) in high and low feed groups. Regression line is for high feed fish only ( $\text{Log}_e G_w = 1.97 - 0.47 \text{Log}_e W_t$ ). (D) Log<sub>e</sub> of specific growth rate of fish of unit size (a comparative measure of growth rate which is independent of body size, see text for explanation), as a function of time for high and low feed groups under normal photoperiod conditions.




Fig. 3. Diel cycles of plasma thyroxine, osmotic and ionic concentrations, and hematocrit (mean  $\pm$  1 standard error of the mean) over a 20 hr period. Only plasma thyroxine, osmolarity and  $[\text{Na}^+]$  has significant diel cycles ( $P < 0.05$ , one-way ANOVA). Experiment was conducted on February 12 under normal photoperiod conditions (10.4 hr daylight, 13.6 hr darkness). The clear horizontal bar indicates daylight period, darkened bar indicates darkness. Sample size was six fish for each time interval, except for the 1200 and 0200 intervals when only four samples for  $[\text{Na}^+]$ ,  $[\text{K}^+]$  and  $[\text{Mg}^{2+}]$  were used.

size, as a function of time (Fig. 2(D)), shows similar growth in the two feeding treatments in March, 1981, corresponding to first feeding. Growth rates from April to December 1981 are much lower for the low feeding group, after which growth in the two groups became more similar, though still lower in the low feed group except for the last weighing. Mean condition factor ( $[\text{weight} \cdot (\text{length}^3)^{-1}] \cdot 100$ ) in each group at every time interval was greater than 0.95 (range: 0.96–1.36, low feed group; 1.09–1.41, high feed group).

Similarity of growth rates in high and low feed groups in March 1981 reflects the equal feeding rates given the two groups just prior to this period. For an 8-month period, smaller ration size in the low feed group drastically reduced growth rate. Similarity of growth rates, from December 1981 on, may reflect a growth-ration relationship which changes with body size though little is known of this function (Ricker, 1979). Reduction in growth rate of high feed fish at large body size may result from a combination of maturation and tank size which may act more strongly in larger fish to inhibit growth.

#### Diel cycles

Plasma  $[\text{Cl}^-]$ ,  $[\text{K}^+]$ ,  $[\text{Mg}^{2+}]$  and hematocrit showed no significant diel cycle ( $P > 0.10$ , ANOVA, Fig. 3). Plasma osmolarity,  $[\text{Na}^+]$  and thyroxine concentration, however, significantly changed over a 20-hr period ( $P < 0.01$ , ANOVA, Fig. 3). Plasma osmolarity peaked after 8.5 hr of light. Both  $[\text{Na}^+]$  and osmolarity declined by the first night sample (Fig. 3). Thyroxine levels were highest during the light period, declined prior to dusk, and reached their lowest levels during darkness. There was no significant difference in variance over the 20-hr period of any of the plasma variables ( $P > 0.10$ , Bartlett's test), indicating that no change in variability occurred as a result of diel cycles.

Diel cycles of plasma ions have been found in other teleosts and are considered to be rhythmic responses to changes in activity or light levels. Hannah and Pickford (1981) found "afternoon" peaks of plasma  $[\text{Na}^+]$  in killifish, *Fundulus heteroclitus*, which did not occur for  $[\text{Cl}^-]$  and  $[\text{K}^+]$ . They also determined a daytime rise in hematocrit, which was not observed in brook trout or juvenile sockeye salmon, *Oncorhynchus nerka* (Leatherland *et al.*, 1974). Although our results show that peaks in  $[\text{Na}^+]$  and total osmolarity were concurrent, increases in "afternoon"  $[\text{Na}^+]$  are not large and cannot account for the more substantial increases in total osmolarity (see Fig. 3). An unmeasured plasma constituent must make up the remaining portion. Hannah and Pickford (1981) hypothesized that "afternoon" increases and "post-sunset" declines in  $[\text{Na}^+]$  may be due to locomotor activity which for killifish and brook trout is high during daylight and low at night. Wood and Randall (1973) have shown that plasma  $[\text{Na}^+]$  increases with activity in rainbow trout. Alternatively, daily  $[\text{Na}^+]$  and osmolarity changes may be related to feeding activity. Although our animals were not fed during the day of sampling, ionic and osmotic cycles may still be due to rhythms associated with daytime feeding. Using an experimental design similar to ours, in which fish were starved overnight, Leatherland *et*

Table 1. Size, age, photoperiod and feeding effects on plasma parameters and gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase. Sample size ( $N$ ), range, mean, standard deviation (SD), and slope ( $b$ ),  $y$ -intercept ( $y$ -int.), coefficient of determination ( $r^2$ ), and significance of regression slope ( $P$ ), are given for each physiological variable and their regression on length and age. Brook trout were 6.0 and 30.5 cm fork length and 180 to 700 d old. Feeding and photoperiod effect were determined using two-way ANOVA (Yes,  $P < 0.05$ ; No,  $P > 0.05$ ). Yes (H) indicates that photoperiod effect was only significant in high (H) feeding group

	$N$	Range	Mean	SD	Independent variable	Linear regression				Feeding effect	Photoperiod effect
						$b$	$y$ -int.	$r^2$	$P$		
Osmolarity	793	272–362	307	12.4	Length	0.64	296	0.12	<0.01	YES	NO
					Age	0.022	298	0.07	<0.01		
[ $\text{Na}^+$ ]	535	126–186	152	8.0	Length	0.33	146	0.06	<0.01	NO	NO
					Age	0.013	147	0.06	<0.01		
[ $\text{K}^+$ ]	607	0.10–7.12	1.99	1.24	Length	-0.037	2.68	0.03	<0.01	YES	NO
					Age	0.0009	2.42	0.01	<0.01		
[ $\text{Mg}^{2+}$ ]	605	0.64–2.23	1.04	0.22	Length	—	—	—	0.47	NO	YES(H)
					Age	—	—	—	0.12		
[ $\text{Cl}^-$ ] (100 l)	352	96–133	108	5.5	Length	0.13	105	0.02	0.02	—	—
					Age	0.012	102	0.08	<0.01		
[ $\text{Cl}^-$ ] (1000 l)	311	103–142	124	6.6	Length	-0.15	127	0.02	0.02	—	—
					Age	-0.008	128	0.03	<0.01		
Hematocrit	940	30–72	48	7.2	Length	0.47	40	0.18	<0.01	—	—
					Age	0.01	44	0.04	<0.01		
$\text{Na}^+$ , $\text{K}^+$ -ATPase	687	1.3–21.2	7.9	3.4	Length	—	—	—	0.52	NO	NO
					Age	—	—	—	0.60		
Thyroxine	728	0.0–10.1	3.01	1.87	Length	0.14	0.48	0.20	<0.01	YES	YES(H)
					Age	0.0045	1.05	0.10	<0.01		

al. (1974) found that plasma free fatty acids of juvenile sockeye salmon peaked and declined during daylight. Other nutrients and waste products may cycle in a similar fashion and result in the observed "afternoon" peak in plasma osmolarity.

Diel cycles of thyroxine, which in the present study peaked during daylight, have been observed in other teleosts. White and Henderson (1977) reported levels of thyroxine ( $T_4$ ) and 3,5,3'-triiodo-L-thyronine ( $T_3$ ) in brook trout that were higher at midday and evening than at dawn. Similar diel patterns in  $T_4$  and possibly  $T_3$ , have been reported for rainbow trout and goldfish (Eales *et al.*, 1981; Spieler and Noeske, 1979). In contrast, Osborn *et al.* (1978) described diel cycling of  $T_4$  and  $T_3$  in rainbow trout, in which lowest values were observed during daylight and highest values at night. Since other investigators have failed to find diurnal variations in plasma thyroxine in rainbow trout (Leatherland *et al.*, 1977; Brown *et al.*, 1978), it seems clear that experimental conditions are involved in diel variations. Eales *et al.* (1981) have shown that starvation for 72 hr eliminates diel variations in  $T_4$ . They also demonstrate that it is not the time of feeding which determines the timing of  $T_4$  and  $T_3$  peaks. From their results it appears that feeding stimulates the diel thyroxine cycle, while some other factor (possibly the light-dark period, or the animals locomotor response to it) acts to synchronize it.

Plasma thyroxine concentrations varied 40% over 24 hr, and approximately 31% over the daytime period in which our sampling for annual cycles occurred. These variations probably did not affect our ability to detect seasonal cycles since sampling occurred during the day when thyroxine levels were highest, and because the seasonal variability (ranging over an order of magnitude) was 3–4 times greater than that of the diel cycle. Changes in the magnitude and timing of diel cycling of thyroxine, however, could vary with season. Meier's (1975) review of circadian prolactin and cortisol rhythms in birds has shown that seasonal changes in diel cycles exist and

possess a regulatory function. The role of diel thyroxine cycles in teleosts has yet to be established.

#### Ontogenetic changes in freshwater

*Plasma osmotic and ionic concentrations.* Range, mean and standard deviation of plasma osmolarity, [ $\text{Na}^+$ ], [ $\text{Cl}^-$ ], [ $\text{K}^+$ ] and [ $\text{Mg}^{2+}$ ] are shown in Table 1. Brook trout used for this analysis were between 6.0–30.5 cm fork length, 6–30 months old and contained both mature and immature individuals. Plasma osmolarity, [ $\text{Na}^+$ ] and [ $\text{K}^+$ ] were significantly correlated with length and age (Table 1), but length and age explained little of the variation in these plasma constituents ( $0.02 < r^2 < 0.12$ , Table 1). Plasma [ $\text{Mg}^{2+}$ ] was not significantly correlated with length or age.

Plasma [ $\text{Cl}^-$ ] was significantly correlated with length and age, but in 100 l tanks the relationship was positive ( $r^2 = 0.02$  and  $0.08$ , respectively) while in 1,000 l tanks it was negative ( $r^2 = 0.02$  and  $0.03$ , respectively). Plasma [ $\text{Cl}^-$ ] of fish kept in 100 l and 1,000 l tanks were significantly different ( $P < 0.01$ , student's  $t$ -test). These differences were observed within 24 hr of transfer from 1,000 to 100 l tanks. pH of water was greater in smaller tanks (6.2–6.4) than in the larger tanks (5.8–6.1), and was possibly due to increased aeration in small tanks. Plasma [ $\text{K}^+$ ] showed inconsistent differences under the two culture conditions (being sometimes higher and sometimes lower in 100 l tanks), while plasma osmolarity, [ $\text{Na}^+$ ], [ $\text{Mg}^{2+}$ ], thyroxine concentration, hematocrit, and gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity showed no significant differences between tanks ( $P > 0.05$ , student's  $t$ -test).

Exposure of fish to lethal and sublethal acidic conditions results in decreased plasma [ $\text{Na}^+$ ] and [ $\text{Cl}^-$ ] (Packer and Dunson, 1970; Neville, 1979; McDonald *et al.*, 1980; Holeyton *et al.*, 1983). Plasma [ $\text{Na}^+$ ] however, did not differ between 1000 l and 100 l tanks. The relatively small differences in pH may account for the absence of plasma [ $\text{Na}^+$ ] reduction. It is interesting to note that while plasma [ $\text{Cl}^-$ ]

of fish in 1001 tanks decreased an average 16 mmole/l, no decrease in total osmolarity occurred, nor was there a substantial change in other plasma ions. An unmeasured plasma constituent, probably anionic, must substitute for decreased plasma  $[\text{Cl}^-]$ .

Feeding regime had a significant effect on plasma osmolarity and  $[\text{K}^+]$ , but not on other plasma ion concentrations (Table 1). When significant differences between high and low feeding groups were observed ( $P < 0.05$ , student's  $t$ -test), plasma osmolarity was always higher in the high feed group, and plasma  $[\text{K}^+]$ , with one exception, was always higher in the low feed group. Significant differences in plasma osmolarity and  $[\text{K}^+]$  between high and low feed groups occurred even for the largest and oldest fish in the low feed group, indicating that meal size itself was exerting an influence.

Photoperiod treatment had a significant effect on plasma  $[\text{Mg}^{2+}]$  (in high feed group only, Table 1), but not on other plasma ions. No consistent difference in  $[\text{Mg}^{2+}]$  due to daylength was found.

Plasma ion and osmotic concentrations of brook trout in freshwater are typical of those reported for other freshwater teleosts (Holmes and Donaldson, 1969). Size and age related adjustments in plasma ion and osmotic concentrations can explain only a small amount of the variation of these parameters. Size related changes in blood ions may not be due to osmoregulatory changes *per se*, but rather to other impinging physiological responses that vary with size. For example, the degree of digestion of the previous days meal is size dependent (Jobling *et al.*, 1977) and could result in nutrient transport related changes in plasma ionic and osmotic concentrations. Despite these alternative explanations, size related changes in hyperosmoregulation may exist.

Significant seasonal changes in plasma ion levels that might signal preparatory physiological adaptations were not found in the present experiment. Seasonal changes in plasma ions have been found in rainbow trout (Lane, 1979; Houston and Smeda, 1979). We found seasonal changes in  $[\text{Mg}^{2+}]$  only,




Fig. 4. Annual variation in hematocrit (% red blood cells) for males (open diamonds) and females (closed diamonds) under normal photoperiod conditions. Mean value of six to ten fish per sample  $\pm 1$  standard error of the mean. Daylength (·) and spawning time (horizontal bar) are also shown. Fish in this experiment were from high feed group, all of which became mature at age 1+ yr in their second autumn (November 1982). Mean female hematocrit in autumn is significantly lower than the non-autumn mean ( $P < 0.05$ , Student's  $t$ -test).

and this effect occurred only in the high feed group. There was no clear trend of variations in plasma  $[\text{Mg}^{2+}]$  with changing daylength in either photoperiod. Lack of photoperiod effect on plasma ions in our experiments, which were conducted at constant temperature, indicates that temperature (or synergy between photoperiod and temperature) may play a more important role than photoperiod alone in determining seasonal ion changes reported by other researchers. Alternatively, brook trout may not display seasonal cycles in ion concentration that are seen in other salmonids.

**Hematocrit.** Hematocrit increases with increasing length which can explain a portion of the variation in hematocrit ( $r^2 = 0.18$ , Table 1), while age can explain little of the variation ( $r^2 = 0.04$ ). This relationship held true when mature fish were excluded ( $r^2 = 0.13$  and 0.09 for length and age, respectively). Adult male and female hematocrits of 1+ year fish are the same for much of the year (Fig. 4) and do not differ until autumn when male hematocrit rises slightly and female hematocrit drops significantly from the spring and summer average ( $P < 0.01$ , Student's  $t$ -test). Changes in hematocrit occur simultaneously with final maturation, when sperm is running freely and egg diameters are at a maximum. Significant male-female differences in hematocrit, however, were also observed in immature fish; immature males and females in autumn photoperiod (11.5–9.5 hr daylength) had mean hematocrits of 50% ( $n = 15$ ) and 44%, respectively ( $n = 13$ ;  $P < 0.01$ , Student's  $t$ -test). Mean hematocrits of immature males and females during winter photoperiod (9.1–10.2 hr daylength) were not significantly different (44% and 46%,  $N = 15$  and 11, respectively,  $P > 0.25$ ).

Hematocrit levels and their variability as reported here are typical of those reported elsewhere for brook trout and other salmonids (Sniezko, 1960; Sano, 1960). Since size explains a greater portion of the variation in hematocrit than age, age is probably significant only to the extent that it covaries with size. Adult male rainbow trout, pike (*Esox lucius*) and largemouth bass (*Micropterus salmoides*) have higher hematocrit than females (Sano, 1960; Mulcahy, 1970; Steuke and Atherton, 1965) indicating that sexual differences in teleost hematocrit are not rare. Sano (1960) also reported a sharp reduction in hematocrit of adult rainbow trout of both sexes that was correlated with gonadal development. In brook trout, only the female hematocrit declines during the onset of spawning. Further work is necessary to determine the mechanistic control of hematocrit and how this control may be related to sex, spawning and photoperiod cycles.

**Gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase.** Individual gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activities ranged from 1.3 to 21.2  $\mu\text{MP}_i \times \text{mg prot.}^{-1} \cdot \text{hr}^{-1}$ , with a mean value of 7.9 (Table 1). Gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity (and its log transformation) were not significantly correlated with size or age. Mean value of gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase did not rise above 13.0  $\mu\text{MP}_i \cdot \text{mg prot.}^{-1} \cdot \text{hr}^{-1}$  for any sampling period. Feeding treatment had no effect on gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase levels in either photoperiod (Table 1), nor did the two photoperiod treatments differ in their effect on gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity.

None of the experimental manipulations of the




Fig. 5. Annual cycles of plasma thyroxine in high feed (squares) and low feed (circles) groups in normal photoperiod (A, solid symbols) and 3 month delayed photoperiod (B, open symbols), as a function of calendar date of sampling. Mean value of 5–16 fish per sample  $\pm$  standard error of the mean. An asterisk (\*) next to mean of high feed group indicates a significantly ( $P < 0.05$ , Student's  $t$ -test) higher mean plasma thyroxine levels than the low feed group at that sampling period. Each point represents samples taken on a single day except for three instances (June, July and August 1981) when samples taken within 2–4 days were combined. Daylength (°) and time of spawning (horizontal bar) are shown for each photoperiod. Feeding treatment had a significant effect on thyroxine levels in both photoperiods ( $P < 0.01$ , two-way ANOVA). Photoperiod treatment significantly affected thyroxine levels in the high feed group ( $P < 0.01$ , two-way ANOVA), but not the low feed group ( $P = 0.43$ ).

present study affected gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity of brook trout. In contrast, gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity in migratory Pacific salmon (*Oncorhynchus* sp.), Atlantic Salmon (*Salmo salar*) and steelhead trout (*Salmo gairdneri*) exhibit a seasonal cycle (usually peaking in spring or autumn, and corresponding to the period of seaward migration) which is synchronized by photoperiod (Zaugg and McLain, 1970; Zaugg and Wagner, 1973; Saunders and Henderson, 1978). Ewing *et al.* (1979) found that although chinook salmon displayed a seasonal cycle of gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity under simulated normal photoperiod conditions, a 3 month advanced photoperiod did not significantly alter the cycle, indicating the controlling influence of endogenous rhythms. In the present study there was no daylength-related rhythm of gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity in either photoperiod treatment, and no evidence of endogenous rhythms.

Ewing *et al.* (1979) found that growth rate of chinook salmon (altered by changes in temperature) affected the cyclic annual change in gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity and that size was positively correlated with this activity. Size, growth and photoperiod did not alter gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity of brook trout in the present study, possibly due to the less specialized nature of its seaward migration. Brook trout, and charr in general, show less anadromy than other salmonids (Hoar, 1976), and there are more non-migratory than migratory populations of brook trout (Power, 1980). Anadromous brook trout spend long periods in estuaries (Montgomery *et al.*, unpublished data) where gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity increases (McCormick and Naiman, 1984b), indicating that adaptations for seawater entry have an important behavioral component.

It is possible that possession of preparatory changes in gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity is genetically determined in brook trout. However, the migratory pattern of the hatchery stock used in these experiments did not differ from that of natural populations (Mullan, 1958). Furthermore, gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activities of an anadromous population of brook trout, which show external signs of smolting (silvering), were not significantly different from a nearby non-anadromous population (McCormick and Naiman, 1984b).

**Plasma thyroxine.** Plasma thyroxine concentration was significantly correlated with both size and age ( $P < 0.01$ ) and explained 20% and 10% of the thyroxine variation, respectively (Table 1). The significance of these correlations may be explained, in part, by changes in growth rate. Within each photoperiod, feeding treatment resulted in significant differences in thyroxine levels (Table 1 and Fig. 5). Fish in high feed groups had significantly higher plasma thyroxine at most sampling times starting from the first sample period (June 1981) and continuing until December 1981. This is the same time that growth rates in the high feed group were much greater than in the low feed group (Fig. 2(D)). After this period, growth rates and thyroxine levels were similar for both feeding groups until April 1982, when high feed fish again attained higher plasma thyroxine levels. This pattern was similar for both photoperiods (Fig. 5(A, B)). Under normal photoperiod conditions the percent mean difference in plasma thyroxine (between the high and low feed group) and the mean difference in specific growth rate per unit body weight (a) 1–2 week later were significantly correlated ( $r = 0.71$ ). These results indicate that higher growth rates in brook trout are associated with higher levels of plasma thyroxine.

Under normal photoperiod conditions (Fig. 5A) there was a strong pattern of high thyroxine levels in the “spring” (increasing photoperiod), low “summer” levels which rose to a secondary “autumn” peak. This pattern was consistent in both high and low feed fish, with the exception that the “spring” peak was attenuated in the low feed group.

Although a single “spring” peak occurred under the 3-month delayed photoperiod for the high feed fish, there was no rise in thyroxine levels in either group during the second “spring”, and there was no

subsequent "summer" decline (Fig. 5B). As such, there was no clearly discernible daylength pattern in thyroxine levels under 3-month delayed photoperiod.

Photoperiod treatment had a significant effect only on fish in the high feed group (Table 1). There was no clear pattern in the differences between normal and delayed photoperiods for the high feed group (Fig. 5(A,B)), and it was clear that a simple 3-month shift in the  $T_4$  cycle did not occur as a result of treatment with a 3-month delayed photoperiod.

The significant effect of feeding level on circulating  $T_4$  concentrations in brook trout is probably related to growth, since significant differences in  $T_4$  levels in high and low feeding groups occurred when growth rates of the two groups were most different. It is unlikely that insufficient iodine in the low feeding group resulted in lower  $T_4$  levels since under normal laboratory conditions less than 20% of the iodine needed by rainbow trout is obtained from the diet (Hunt and Eales, 1979), and only 5% of the total iodine is used by the thyroid. In addition, brook trout deprived of food for several weeks increase their plasma iodine (Higgs and Eales, 1971), while rainbow trout show no detectable change in  $T_4$  after up to 40 days of starvation (Leatherland *et al.*, 1977; Milne *et al.*, 1979). More recent experiments indicate that drastically reduced feeding may decrease plasma  $T_4$  of salmonids (Dickhoff, unpublished; Higgs *et al.*, 1982). Direct feeding effects cannot accurately describe our results, however, since feeding treatment lasted throughout the study while differences in  $T_4$  between high and low feed groups occurred for a limited period. Thyroid hormones, particularly triiodothyronine, administered exogenously can stimulate growth in a variety of teleosts and most salmonids (Higgs *et al.*, 1982).  $T_4$  may act in synergy with other anabolic hormones, particularly growth hormone, to stimulate somatic growth (Donaldson *et al.*, 1979). Our findings of increased circulating  $T_4$  levels associated with higher growth rates support a model of thyroid influence on growth.

A seasonal pattern of circulating  $T_4$ , similar to that reported here was found in adult brook trout by White and Henderson (1977), with the one exception that a secondary fall peak was not found. This seasonal pattern in brook trout  $T_4$  levels is similar to that found for smolting salmonids (Dickhoff *et al.*, 1982). The magnitude of the springtime peak, however, is generally greater in smolting salmonids. The lack of preparatory seawater-entry adaptations in brook trout (such as increases in gill  $Na^+$ ,  $K^+$ -ATPase) suggests that spring thyroxine increases activate other physiological functions. The thyroxine cycle displayed in primitive salmonids such as brook trout has perhaps been sequestered by specialized migrators to synchronize migration and smoltification.

Winter flounder (*Pseudopleuronectes americanus*) displayed peak thyroxine concentrations in spring and low concentrations in autumn (Eales and Fletcher, 1982), while high winter concentrations and low summer concentrations were observed in rainbow trout (Osborn *et al.*, 1978). Constant temperatures used in our experiments indicate that temperature changes are not necessary to elicit seasonal thyroxine cycles. Although feeding activity increased

during spring photoperiod in our experiment (personal observation), we did not detect increased growth rates under increasing photoperiod and therefore cannot ascribe higher spring thyroxine levels to increased growth during this period.

The delayed photoperiod regime did not shift thyroxine cycles 3 months from the normal regime; in low feed fish photoperiod has no effect on plasma thyroxine, while in high feed fish the effect seemed to be a dampening of the normal cycle. These results raise the possibility that an endogenous cycle, or an exogenous cycle synchronized by an environmental factor other than photoperiod or temperature, exists in brook trout. Other photoperiod cued cycles, in particular maturation, did respond to the photoperiod treatment; final maturation under our experimental conditions was delayed 3 months in the delayed photoperiod (McCormick and Naiman, 1984c). These results are somewhat conflicting, especially in light of other evidence associating thyroid changes with maturation (see Leatherland, 1982, for review). Nonetheless, it appears that an annual cycle of thyroxine with a spring peak is not necessary to begin or synchronize the maturation cycle of brook trout.

#### SUMMARY

Significant diel cycles were observed in plasma osmolarity,  $[Na^+]$  and thyroxine concentration, and were not detected in plasma  $[Cl^-]$ ,  $[K^+]$ ,  $[Mg^{2+}]$  and hematocrit. Plasma osmolarity,  $[Na^+]$  and thyroxine concentrations were highest during daylight and lowest at night. This cycle may be caused by feeding and locomotor activity which are highest during periods of light.

Plasma osmolarity,  $[Na^+]$ ,  $[K^+]$  and hematocrit increased with increasing size and/or age of brook trout, and can explain a small portion of their variation. Plasma osmolarity and  $[K^+]$  were also influenced by feeding level. The effect of size on plasma ions may be explained by a more favorable surface area to volume ratio which, other things being equal, will result in lower net water influx, lower plasma water, higher plasma ions and higher hematocrit with increasing size. We cannot, however, rule out other factors which may also covary with size and/or age.

Gill  $Na^+$ ,  $K^+$ -ATPase activity in brook trout did not respond to feeding or photoperiod treatment, nor was there evidence of size, age or daylength related changes. Brook trout therefore do not possess preparatory physiological adaptations in gill  $Na^+$ ,  $K^+$ -ATPase that are characteristic of other migratory salmonids. It would appear that the more variable and opportunistic nature of brook trout migrations has not resulted in sufficient selection pressure for the development of preparatory, photoperiod-controlled changes in gill  $Na^+$ ,  $K^+$ -ATPase.

Plasma thyroxine concentrations were higher in high feed fish and were directly correlated with size. Significant differences in plasma thyroxine concentration between high and low feed groups occurred when differences in growth between the two groups were greatest. These results are best explained by an



interaction between growth rate and plasma thyroxine. Under normal photoperiod conditions, plasma thyroxine exhibited a seasonal cycle consisting of high levels in spring, low summer levels and a secondary peak in autumn. Three-month delayed photoperiod did not result in a shift of the thyroxine cycle. Since seasonal changes of brook trout gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity and hypoosmoregulatory ability did not occur (McCormick and Naiman, 1984a), the annual cycle of plasma thyroxine does not stimulate these physiological functions as it is presumed to in smolting salmonids. The seasonal thyroxine cycle which exists in the more primitive charrs must exert its influence through other seasonally occurring physiological functions such as growth, activity or appetite.

*Acknowledgements*—We thank the Massachusetts Division of Fisheries and Wildlife for providing brook trout eggs. M. Zaltas, M. Connor, E. Rosa-Molinar, R. Rheault and E. Montgomery ably provided technical assistance. V. Marsard typed the first draft. J. M. Capuzzo, R. B. Gagosian, R. W. Griffith, J. J. Stegeman and G. C. Walker gave helpful advice during the conduct of our experiments and made many constructive comments in review. This project was supported by the U.S. Department of Commerce, NOAA, Office of Sea Grant NA80-AA-D-00077 (R/A-14). S. D. McC. was supported by a Tai-Ping Predoctoral Fellowship in Marine Biology and the Woods Hole Oceanographic Institution Education Office. This is Contribution No. 559 of the Woods Hole Oceanographic Institution and Contribution No. 85 of the Institution's Matamek Research Station.

#### REFERENCES

- Brett J. R. (1979) Environmental factors and growth. In: *Fish Physiology, Vol. VIII* (edited by Hoar W. S., Randall D. J. and Brett J. R.), pp. 599–675. Academic Press, New York.
- Brown J., Fedoruk K. and Eales J. G. (1978) Physical injury due to injections or blood removal causes elevations of plasma thyroxine in rainbow trout, *Salmo gairdneri*. *Can. J. Zool.* **56**, 1998–2003.
- Castonguay M., Fitzgerald G. J. and Côté Y. (1982) Life history and movements of anadromous brook charr, *Salvelinus fontinalis*, in the St. Jean River, Gaspé, Québec. *Can. J. Zool.* **60**, 3084–3091.
- Dickhoff W. W., Darling D. S. and Gorbman A. (1982) Thyroid function during smoltification of salmonid fish. In: *Gunma Symposia on Endocrinology, Vol. 19, Phylogenetic Aspects of Thyroid Hormone Actions*. (edited by Institute of Endocrinology, Gunma Univ.), pp. 45–61. Center for Academic Publications Japan, Tokyo.
- Dickhoff W. W., Folmar L. C. and Gorbman A. (1978) Changes in plasma thyroxine during smoltification of coho salmon, *Oncorhynchus kisutch*. *Gen. Comp. Endocrinol.* **36**, 229–232.
- Donaldson E. M., Fagerlund U. H. M., Higgs D. A. and McBride J. R. (1979) Hormonal enhancement of growth in fish. In: *Fish Physiology, Vol. VIII*. (edited by Hoar W. S., Randall D. J. and Brett J. R.), pp. 456–597. Academic Press, New York.
- Dutil J. D. and Power G. (1980) Coastal populations of brook trout, *Salvelinus fontinalis*, in Lac Guillaume-Delisle (Richmond Gulf) Québec. *Can. J. Zool.* **58**(10), 1828–1835.
- Eales J. G. and Fletcher G. L. (1982) Circannual cycles of thyroid hormones in plasma of winter flounder (*Pseudopleuronectes americanus* Walbaum). *Can. J. Zool.* **60**, 304–309.
- Eales J. G., Hughes M. and Uin L. (1981) Effect of food intake on diel variation in plasma thyroid hormone levels in rainbow trout, *Salmo gairdneri*. *Gen. Comp. Endocrinol.* **45**, 167–174.
- Ewing R. D., Johnson S. L., Pribble H. J. and Lichatowich J. A. (1979) Temperature and photoperiod effects on gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity in chinook salmon (*Oncorhynchus tshawytscha*). *J. Fish. Res. Bd Can.* **36**, 1347–1353.
- Gorbman A., Dickhoff W. W., Mighell J. L., Prentice E. F. and Waknitz F. W. (1982) Morphological indices of developmental progress in the parr-smolt coho salmon, *Oncorhynchus kisutch*. *Aquaculture* **28**, 1–19.
- Grau E. G., Specker J. L., Nishioka R. S. and Bern H. A. (1982) Factors determining the occurrence of the surge in thyroid activity during smoltification. *Aquaculture* **28**, 49–57.
- Hannah S. G. and Pickford G. E. (1981) Diurnal peaks in hematocrit and in serum sodium in the killifish, *Fundulus heteroclitus*, and their absence in serum potassium and chloride: a lack of correlation. *Comp. Biochem. Physiol.* **70A**, 157–159.
- Higgs D. A. and Eales J. G. (1971) Iodide and thyroxine metabolism in the brook trout, *Salvelinus fontinalis* (Mitchell) during sustained exercise. *Can. J. Zool.* **49**, 1255–1269.
- Higgs D. A., Fagerlund U. H. M., Eales J. G. and McBride J. R. (1982) Application of thyroid and steroid hormones as anabolic agents in fish culture. *Comp. Biochem. Physiol.* **73B**, 143–176.
- Hoar W. S. (1976) Smolt transformation: evolution, behavior and physiology. *J. Fish Res. Bd Can.* **33**, 1234–1252.
- Holeton G. F., Neumann P. and Heisler N. (1983) Branchial ion exchange and acid-base regulation after strenuous exercise in rainbow trout (*Salmo gairdneri*). *Resp. Physiol.* **51**, 303–318.
- Holmes W. N. and Donaldson E. M. (1969) The body compartments and the distributions of electrolytes. In: *Fish Physiology, Vol. I* (edited by Hoar W. S. and Randall, D. J.), pp. 1–90. Academic Press, New York.
- Houston A. H. and Smeda P. S. (1979) Thermoacclimatory changes in the ionic microenvironment of hemoglobin in the stenothermal rainbow trout (*Salmo gairdneri*) and eurythermal carp (*Cyprinus carpio*). *J. exp. Biol.* **80**, 317–340.
- Hunt D. W. C. and Eales J. G. (1979) Iodide balance in rainbow trout, *Salmo gairdneri* and effects of testosterone propionate. *J. Fish. Res. Bd Can.* **36**, 282–285.
- Jobling M. (1983) Growth studies with fish—overcoming the problems of size variation. *J. Fish Biol.* **22**, 153–157.
- Jobling M., Gwyther D. and Grove D. J. (1977) Some effects of temperature, meal size and body weight on gastric evacuation time in the dab, *Limanda limanda* (L.). *J. Fish Biol.* **10**, 291–298.
- Lane H. C. (1979) Progressive changes in hematology and tissue water of sexually mature trout (*Salmo gairdneri*) Richardson, during autumn and winter. *J. Fish Biol.* **15**, 425–436.
- Leatherland J. F. (1982) Environmental physiology of the teleostean thyroid gland: a review. *Env. Biol. Fish.* **7**, 83–110.
- Leatherland J. F., McKeown B. A. and John T. M. (1974) Circadian rhythm of plasma prolactin, growth hormone, glucose and free fatty acid in juvenile kokanee salmon, *Oncorhynchus nerka*. *Comp. Biochem. Physiol.* **74A**, 821–828.
- Leatherland J. F., Cho C. Y. and Slinger S. J. (1977) Effects of diet, ambient temperature and holding conditions on plasma thyroxine levels in rainbow trout (*Salmo gairdneri*). *J. Fish. Res. Bd Can.* **34**, 677–682.
- Leitritz E. and Lewis R. C. (1976) Trout and salmon culture. *Calif. Fish Game Bull.* **164**, pp. 220.
- Lowry O. H., Roseborough N. J., Farr A. L. and Randall

- R. J. (1951) Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**, 265-275.
- McCormick S. D. and Naiman R. J. (1984a) Osmoregulation in the brook trout, *Salvelinus fontinalis*. II. Effects of size, age and photoperiod on seawater survival and ionic regulation. *Comp. Biochem. Physiol.* **79A**.
- McCormick S. D. and Naiman R. J. (1984b) The Physiology of Smoltification in Anadromous and Non-anadromous Brook Trout (*Salvelinus fontinalis*) and Atlantic Salmon (*Salmo salar*) from the Matamek River and Rivière à la Truite, Québec. *Can. J. Fish. Aqu. Sci.* (in press).
- McCormick S. D. and Naiman R. J. (1984c) Some determinants of maturation in the brook trout, *Salvelinus fontinalis*. *Aquaculture* (in press).
- McDonald D. G., Hobe H. and Wood C. M. (1980) The influence of calcium on physiological responses of the rainbow trout, *Salmo gairdneri*, to low environmental pH. *J. exp. Biol.* **88**, 259-267.
- Meier A. H. (1975) Chronoendocrinology of vertebrates. In: *Hormonal Correlates of Behavior*, Vol. 2. (edited by Eleftherious, B. E. & Sprout, R. L.), pp. 469-549. Plenum Press, New York.
- Miller G. L. (1959) Protein determination for large numbers of samples. *Anal. Chem.* **31**, 964.
- Milne R. S., Leatherland J. F. and Holub B. J. (1979) Changes in plasma thyroxine, triiodothyronine and cortisol associated with starvation in rainbow trout (*Salmo gairdneri*). *Env. Biol. Fish* **4**, 185-190.
- Mulcahy M. F. (1970) Blood values in the pike *Esox lucius* L. *J. Fish Biol.* **2**, 203-209.
- Mullan J. W. (1958) The sea-run or "salter" brook trout (*Salvelinus fontinalis*) fishery of the coastal streams of Cape Cod, Massachusetts. *Mass. Div. Fish and Game. Bull. No. 17*.
- Neville C. M. (1979) Influence of mild hypercapnia on the effects of environmental acidification on rainbow trout (*Salmo gairdneri*). *J. exp. Biol.* **83**, 345-349.
- Osborn R. H., Simpson T. H. and Youngson A. F. (1978) Seasonal and diurnal rhythms of thyroidal status in the rainbow trout, *Salmo gairdneri* Richardson. *J. Fish Biol.* **12**, 531-540.
- Packer R. K. and Dunson W. A. (1970) Effects of low environmental pH on blood pH and sodium in brook trout. *J. exp. Zool.* **174**, 65-71.
- Power G. (1980) The brook charr, *Salvelinus fontinalis*. In: *Charrs* (edited by Balon, E. K.), pp. 141-203. Dr. W. Junk by Publishers, The Hague.
- Ricker W. E. (1979) Growth rates and models. In: *Fish Physiology*, Vol. VIII (edited by Hoar, W. S., Randall, D. J. & Brett, J. R.), pp. 678-737. Academic Press, New York.
- Sano T. (1960) Haematological studies of the culture fishes in Japan. 3. Changes in blood constituents with growth of rainbow trout. *J. Tokyo Univ. Fisheries* **46**, 77-87.
- Saunders R. L. and Henderson E. B. (1978) Changes in gill ATPase activity and smolt status of Atlantic salmon (*Salmo salar*). *J. Fish. Res. Bd. Can.* **35**, 1542-1546.
- Snieszko S. F. (1960) Microhematocrit as a tool in fishery research and management. *U.S. Fish and Wildl. Serv. Spec. Rep. Fish.* **341**, 1-15.
- Spieler R. E. and Noeske T. A. (1979) Diel variations in circulating levels of triiodothyronine and thyroxine in goldfish, *Carrassius auratus*. *Can. J. Zool.* **57**, 665-669.
- Steucke E. E. and Atherton C. R. (1965) Use of microhaematocrit values to sex largemouth bass. *Prog. Fish Cult.* **27**, 87-89.
- Sutterlin A. M., Harmon P. and Barchard H. (1976). The culture of brook trout in salt water. *Can. Fish. Mar. Serv. Tech. Rept. No. 636*.
- Wedemeyer G. A., Saunders R. L. and Clarke W. C. (1980) Environmental factors affecting smoltification and early marine survival of anadromous salmonids. *Mar. Fish. Rev.* **46**, 1-14.
- White H. C. (1940) Life-history of sea-running brook trout (*Salvelinus fontinalis*) of Moser river, Nova Scotia. *J. Fish. Res. Bd. Can.* **5**, 176-186.
- White B. A. and Henderson H. E. (1977). Annual variation in the circulating levels of thyroid hormones in the brook trout, *Salvelinus fontinalis*, as measured by radioimmunosassay. *Can. J. Zool.* **55**, 475-481.
- Whoriskey F. G., Naiman R. J. and Montgomery W. L. (1981) Experimental sea ranching of brook trout, *Salvelinus fontinalis*, Mitchill. *J. Fish Biol.* **19**, 637-651.
- Wilder D. G. (1952) A comparative study of anadromous and freshwater populations of brook trout (*Salvelinus fontinalis* (Mitchill)). *J. Fish. Res. Bd. Can.* **9**, 169-203.
- Wood C. M. and Randall D. J. (1973) The influence of swimming activity on sodium balance in the rainbow trout (*Salmo gairdneri*). *J. Comp. Physiol.* **82**, 207-233.
- Zaugg W. S. (1982) A simplified preparation for adenosine triphosphatase determination in gill tissue. *Can. J. Fish. Aqu. Sci.* **39**, 215-217.
- Zaugg W. S. and McLain L. R. (1970) Adenosine triphosphatase activity in gills of salmonids: seasonal variation and salt water influence in coho salmon, *Oncorhynchus kisutch*. *Comp. Biochem. Physiol.* **35**, 587-596.
- Zaugg W. S. and Wagner H. H. (1973) Gill ATPase activity related to parr-smolt transformation and migration in steelhead trout (*Salmo gairdneri*): influence of photoperiod and temperature. *Comp. Biochem. Physiol.* **45B**, 955-965.